THE PRESENCE OF PUTATIVE VIRULENCE DETERMINANTS, TETRACYCLINE AND β - LACTAMS RESISTANCE GENES OF *Aeromonas* SPECIES ISOLATED FROM PET TURTLES AND THEIR ENVIRONMENT

S.H.M.P Wimalasena, Gang-Joon Heo*

Laboratory of Aquatic Animal Medicine, Veterinary Medical Center and College of Veterinary Medicine, Chungbuk National University, Chungdae-ro 1, Seowon-gu, Cheongju 28644, Republic of Korea

*Corresponding author, E-mail: gjheo@cbu.ac.kr

Abstract: This study aimed to characterize *Aeromonas* spp. isolated from ten popular species of pet turtles and their environment to evaluate the potential risk of pet turtles as a source of virulence-associated genes, and tetracycline and β -lactams resistance determinants. Presence of eight virulence genes (*ser*, *aer*, *exu*, *lip*, *fla*, *ascV*, *ahyB* and *gcaT*), and tetracycline (*tetA*, *tetB* and *tetE*) and β -lactams (*bla*_{TEM}, *bla*_{SHV}, *bla*_{OXA} and *bla*_{CTX-M}) resistance genes were evaluated by conventional PCR assays. The *aerA* gene showed the highest frequency of occurrence (92%), followed by *fla* (75%), *gcaT* (68%), *ahyB* (59%), *ser* (39%), *lip* (37%) and *ascV* (25%) genes. None of the isolates carried amplicon of DNase-associated *exu* gene. *A. hydrophila*, *A. dharmensis*, *A. veronii* and *A. caviae* were carried seven tested virulence genes except for *exu* while only four virulence genes were detected in *A. enteropelogenes*. Among the 75 tetracycline-resistant isolates, *tetA*, *tetE* and *tetB* genes were detected in 38, 26 and 6 isolates, respectively. Among the tested β -lactam resistance genes, *bla*_{OXA} and *bla*_{TEM} genes were detected in 54% and 36% of β -lactam resistant isolates, respectively. No *bla*_{CTX-M} and *bla*_{SHV} genes were detected. Our results indicate that pet turtle-associated aeromonads, exhibiting potential virulence and antimicrobial (tetracycline and β -lactams) resistance genes, may pose a serious health risk to pet turtle owners, particularly to immunocompromised individuals.

Key words: *Aeromonas* spp.; virulence-associated genes; tetracycline resistance; β -lactams resistance; pet turtle

Introduction

Mesophilic aeromonads are ubiquitous bacteria that are a component of the normal microbiota of many aquatic animals such as fish, amphibians, and reptiles (1). They can cause ulcerative stomatitis, pneumonia, dermatitis, and septicemia in reptiles under stressful conditions such as trapping, handling and temperature variations of rearing environment (2, 3). Over the years, many studies have been investigated to evaluate the prevalence of *Aeromonas* species in aquatic animals, mainly food-producing animals (4, 5). However, a limited number of

studies evaluating the distribution of aeromonads in pet turtles have been published up to date (6, 7).

The pathogenesis of *Aeromonas* species involves various virulence factors including cytotoxic heat-labile enterotoxin (*act*), cytotoxic heat-labile enterotoxin (*alt*) and cytotoxic heat-stable enterotoxin (*ast*), aerolysin (*aer*), lipase (*lip*), serine protease (*ser*), elastase (*ahyB*), DNase (*exu*), glycerophospholipid-cholesterol acyltransferase (*gcaT*), flagellar system (*fla*) and Type III secretion system (TTSS) effector (*ascV*). These genes encoding virulence factors have been broadly used in determining the potential pathogenicity of *Aeromonas* species isolated from the environment, foodstuffs, and human clinical samples (1, 8-10).

Recently, antibiotic-resistant aeromonads have been recognized as a serious concern due to their potential health risks to animals and humans (11, 12). Especially, the dissemination of tetracycline and β -lactams resistance aeromonads in the aquatic environment has been widely documented (12, 13, 14). Among many tetracycline resistance genes, the *tetE*, *tetA* and *tetB* genes were frequently identified from *Aeromonas* species in the aquatic environment (12, 15, 16). *Aeromonas* species can produce numerous β -lactamases for conferring resistance to β -lactams. According to isolation sources, the previous studies have shown the different prevalence of genes encoding β -lactamases in *Aeromonas* species. In the aquatic environment, the *bla*_{TEM}, *bla*_{SHV}, *bla*_{OXA} and *bla*_{CTX-M} β -lactams genes were frequently detected from *Aeromonas* species (17-19).

These resistance genes containing plasmids and transposons are known as mobile genetic elements that can be transferred horizontally among distantly related lineages. Particularly, The aquatic environment is more favorable for the transmission of resistant bacteria, thus, *Aeromonas* species as opportunistic pathogens might be dangerous vectors for the spreading of antibiotic resistance genes through the aquatic environment (18, 20). Hence, the present study was conducted to determine the occurrence of antimicrobial resistance genes (tetracyclines and β -lactams) and virulence-associated genes of *Aeromonas* species isolated from pet turtles and their environment.

Materials and methods

Bacterial isolates

One hundred and two *Aeromonas* species isolates obtained from ten commercially popular pet turtles species (Chinese stripe-necked turtles *Ocadia sinensis*, yellow belly sliders *Trachemys scripta scripta*, river cooters *Pseudemys concinna concinna*, northern Chinese softshell turtles *Pelodiscus maackii*, western painted turtles *Chrysemys picta bellii*, peninsula cooters *Pseudemys peninsularis*, African sideneck turtles *Pelusios castaneus*, common musk turtles *Sternotherus odoratus*, red belly cooters *Pseudemys rubriventris* and alligator snapping turtles *Macrolemys Temminckii*) and their rearing environment was screened to investigate the presence of putative virulence, and β -lactams and tetracycline resistance genes. These iso-

lates have been previously characterized for their antimicrobial susceptibilities, enterotoxin (*act*, *alt* and *ast*) genes and quinolone resistance determinants (7, 21).

Detection of antibiotic resistance genes

Twenty-eight and seventy-five isolates were selected (21) for the detection of β -lactams and tetracycline resistance determinants, respectively. These isolates were tested by PCR assays to detect the genetic determinants associated with resistance to β -lactams (*bla*_{TEM}, *bla*_{SHV}, *bla*_{OXA} and *bla*_{CTX-M}), and tetracyclines (*tetA*, *tetB* and *tetE*). The primer sets used in PCR amplification are summarized in table 1. PCR amplifications were conducted in 20 μ L volumes consisting of 10 μ L of Quick Taq® HS DyeMix (Toyobo, Japan), 1 μ L of 10 pmol/ μ L each primer and 1 μ L of the template under standard conditions. The PCR products were analyzed by electrophoresis on 2% (wt/vol) agarose gels. Positive controls were implemented with previously characterized enterobacterial strains that harbored the corresponding genes (21, 22).

Detection of virulence-associated genes

All isolates were subjected to PCR assays to detect the 8 tested virulence genes including *ser*, *aer*, *exu*, *lip*, *fla*, *ascV*, *ahyB* and *gcat*. The PCR amplification of the virulence-associated genes was carried out according to the PCR primers and conditions reported previously (Table 1). The PCR mixture of 20 μ L contained 10 μ L Quick Taq HS DyeMix (Toyobo, Japan), 7 μ L PCR water, 1 μ L template and 1 μ L of each primer. The PCR products were examined by electrophoresis on 1.5% (W/V) agarose gel.

Results

Bacterial isolates

One hundred and two *Aeromonas* species isolates were isolated from the feces, skin and rearing environments of pet turtles and identified by biochemical and *gyrB* sequence analyses. *Aeromonas enteropelogenes* was the predominant species among the isolates (52.9%) followed by *A. hydrophila* (32.4%), *A. dhakensis* (5.9%), *A. veronii* (4.9%) and *A. caviae* (3.9%)⁷.

Presence of resistance genes

Among the tested β -lactam resistance genes, *bla*_{OXA} and *bla*_{TEM} genes were detected in 54% and 36% of β -lactam resistant isolates, respectively. No

*bla*_{CTX-M} and *bla*_{SHV} genes were detected (Table 2). Among the 75 tetracycline-resistant isolates, *tetA*, *tetE* and *tetB* genes were detected in 38, 26 and 6 isolates, respectively (Table 3).

Table 1: Oligonucleotide primers and PCR conditions ^a used to amplify virulence and antibiotic resistance genes of *Aeromonas* spp.

Gene	Target	Nucleotide Sequence (5'-3')	Size (bp)	Annealing temperature (°C)	Reference
<i>aerA</i>	Aerolysin	F: CTATGGCCTGAGCGAGAAG R: CAGTTCAGTCCCACCACT	431	62	30
<i>ser</i>	Serine protease	F: ACCGAAGTATTGGGTCAGG R: GCTCATGCGTAACTCTGGT	350	55	13
<i>fla</i>	Flagella	F: CCAACCGTYTGACCTC R: MYTGGTTGCGRATGGT	608	56	36
<i>ahyB</i>	Elastase	F: CACGGTCAAGGAGATCAAC R: GCTGGTGTGGCCAGCAGG	513	58	13
<i>lip</i>	Lipase	F: ATCTTCTCCGACTGGTTTCGG R: CCGTGCCAGGACTGGGTCTT	382	62	36
<i>exu</i>	DNase	F: AGACATGCACAACCTCTTCC R: GATTGGTATTGCCTTGCAAG	323	59	13
<i>gcaT</i>	Glycerophospholipid-cholesterol acyltransferase	F: TCCTGGAATCCCAAGTATCAG R: GCAGGTTGAACAGCAGTATCT	237	65	13
<i>ascV</i>	Type III Secretion System	F: AGCAGATGAGTATCGACGG R: AGGCATTCTCCTGTACCAG	891	58	38
<i>bla</i> _{TEM}	β - lactams resistance	F: ATAAATTCTTGAAGACGAAA R: GACAGTTACCAATGCTTAATC	1080	60	22
<i>bla</i> _{SHV}		F: TTATCTCCCTGTTAGCCACC R: GATTTGCTGATTTTCGCTCGG	795	60	
<i>bla</i> _{CTX-M}		F: CGCTTTGCGATGTGCAG R: ACCGCGATATCGTTGGT	550	52	
<i>bla</i> _{OXA}		F: TCAACTTTCAAGATCGCA R: GTGTGTTTAGAATGGTGA	591	60	
<i>tetA</i>		F: GTAATTCTGAGCACTGTGCG R: CTGCCTGGACAACATTGCTT	1000	62	
<i>tetB</i>	Tetracycline resistance	F: CTCAGTATTCCAAGCCTTTG R: CTAAGCACTTGTCTCCTGTT	400	57	22
<i>tetE</i>		F: GTGATGATGGCACTGGTCAT R: CTCTGCTGTACATCGCTCTT	1100	62	

^a PCR thermocycle conditions for each reaction; initial denaturation of 94 °C for 2 min followed by a total of 35 cycles of amplification. Each cycle consisted of 94 °C denaturation for 30 s, annealing for 50 s and 72 °C extension for 10 min.

Table 2: β -lactams resistance profiles of turtle-associated *Aeromonas* spp.

Isolate	Host ^a	β -lactam resistance ^b	β -lactam resistance genes
<i>Aeromonas caviae</i>			
AD14	CSN	AMP, AMX, CEP, FOX	<i>bla</i> _{TEM}
AC50	RC	AMP, AMX, CEP, CRO, FOX, IMI	<i>bla</i> _{OXA} ² <i>bla</i> _{TEM}
<i>A. dharkensis</i>			
AD17	RC	AMP, AMX, CEP, CRO, FOX, CTX	<i>bla</i> _{OXA} ² <i>bla</i> _{TEM}
AD18	RC	AMP, AMX, CEP, CRO, FOX, CTX	<i>bla</i> _{OXA} ² <i>bla</i> _{TEM}
AD19	RC	AMP, AMX, CEP, CRO, FOX	<i>bla</i> _{OXA}
AD15	CSN	AMP, AMX, CEP, FOX, CTX	<i>bla</i> _{OXA}
<i>A. enteropelogenes</i>			
AC2	RC	AMP, AMX, CEP	<i>bla</i> _{OXA}
AC6	RC	CEP, FOX	-
AC15	NCS	AMP, AMX, CEP, FOX	<i>bla</i> _{OXA}
AC30	CM	CEP, CTX, ATM	-
AC31	WP	CEP, CTX, ATM	-
AC32	WP	CEP, CTX, ATM	-
AC35	RC	CEP, CTX, ATM	-
AC44	YB	AMP, AMX, CRO	<i>bla</i> _{TEM}
AC45	RC	CEP, CRO, ATM	-
AC53	WP	AMP, AMX, CEP, FOX	<i>bla</i> _{OXA}
AV4	RC	AMP, AMX, CEP, FOX	<i>bla</i> _{OXA}
AD1	CM	AMP, AMX, CEP, FOX	<i>bla</i> _{OXA}
<i>A. hydrophila</i>			
AH1	RC	AMP, CEP	-
AH11	CSN	AMP, AMX, CEP, CRO	<i>bla</i> _{OXA} ² <i>bla</i> _{TEM}
AH13	CSN	CEP, CRO, FOX, IMI	-
AH19	NCS	AMP, AMX, CEP, CRO	<i>bla</i> _{OXA}
AH20	NCS	AMP, AMX, CEP	<i>bla</i> _{TEM}
AH22	YB	AMP, AMX, CEP	-
AH23	YB	AMP, AMX, CEP, CRO	<i>bla</i> _{OXA}
AH25	CM	AMP, AMX, CEP, FOX	<i>bla</i> _{TEM}
AD10	AF	AMP, AMX, CEP, FOX	<i>bla</i> _{OXA} ² <i>bla</i> _{TEM}
<i>A. veronii</i>			
AC52	SN	AMP, AMX, CEP, FOX	<i>bla</i> _{OXA} ² <i>bla</i> _{TEM}

^a**Host:** CSN= Chinese stripe-necked turtle, YB= yellow belly slider, RC= river cooter, PC= peninsula cooter, NCS= northern Chinese softshell turtle, CM= common musk turtle, WP= western painted turtle, AF= African sideneck turtle, SN= Alligator snapping turtle.

^b **β -lactams resistance:** AMX=Amoxicillin (10 μ g), AMP=Ampicillin (10 μ g), CEP=Cephalothin (30 μ g), CRO=Ceftriaxone (30 μ g), FOX=Cefoxitin (30 μ g), CTX=Cefotaxime (30 μ g), IMI=Imipenem (10 μ g)

Table 3: Distribution of tetracycline resistance genes among tetracycline resistant *Aeromonas* species isolated from pet turtles and their environment

Species	Number of positive isolates (Subtotal %)		
	<i>tetA</i>	<i>tetB</i>	<i>tetE</i>
<i>Aeromonas enteropelogenes</i> (n = 50)	32 (64)	-	8 (2)
<i>A. hydrophila</i> (n = 17)	6 (35)	-	12 (71)
<i>A. dharkensis</i> (n = 4)	-	2 (50)	3 (75)
<i>A. veronii</i> (n = 3)	-	3 (100)	2 (66)
<i>A. caviae</i> (n = 1)	-	1 (100)	1 (100)
Total (%) (n = 75)	38 (51)	6 (1)	26 (35)

Table 4: Prevalence of virulence-associated genes in *Aeromonas* species isolates from pet turtles and their environment

Species	Number of positive isolates (Subtotal %)							
	<i>aerA</i>	<i>lip</i>	<i>ahyB</i>	<i>ser</i>	<i>exu</i>	<i>fla</i>	<i>gcat</i>	<i>ascV</i>
<i>Aeromonas enteropelogenes</i> (n = 54)	46 (85)	0	21 (39)	0	0	51 (94)	23 (43)	0
<i>A. hydrophila</i> (n = 33)	33 (100)	30 (91)	28 (85)	31 (94)	0	19 (58)	33 (100)	15 (47)
<i>A. dharkensis</i> (n = 6)	5 (83)	3 (50)	4 (67)	4 (67)	0	2 (33)	6 (100)	4 (67)
<i>A. veronii</i> (n = 5)	5 (100)	3 (60)	5 (100)	3 (60)	0	2 (40)	4 (80)	3 (60)
<i>A. caviae</i> (n = 4)	4 (100)	2 (50)	2 (50)	2 (50)	0	3 (75)	3 (75)	4 (100)
Total (%) (n = 102)	93 (92)	38 (37)	60 (59)	40 (39)	-	77 (75)	69 (68)	26 (25)

Distribution of virulence-associated genes

The occurrence and frequencies of virulence genes are shown in Table 4. The *aerA* gene showed the highest frequency of occurrence (92%), followed by *fla* (75%), *gcat* (68%), *ahyB* (59%), *ser* (39%), *lip* (37%) and *ascV* (25%) genes. None of the isolates carried amplicon of the DNase-associated *exu* gene.

Discussion

The *Aeromonas* spp. under study were multi-drug-resistant turtle-associated bacteria which carried quinolone resistance determinants, as well as enterotoxin genes (7, 21). The isolates were highly resistant to β -lactams especially amoxicillin, ampicillin and cephalothin. β -lactam antibiotics have been used for the treatment of *Aeromonas* infection during the last decade. However, their efficacy has significantly declined due to the production of β -lactamases by resistant bacterial strains (14, 17, 23). The *Aeromonas* spp. are naturally resistant to β -lactams because of the expression of chromosomal β -lactamases (24).

In this study, twenty-eight aeromonads isolates were resistant to the more than one β -lactam antibiotics. Among them, 54% and 36% of isolates harbored *bla*_{OXA} and *bla*_{TEM} genes. Several previous studies have documented the detection of the *bla*_{OXA} and *bla*_{TEM} genes in *Aeromonas* isolates recovered from the environment (14, 25) and clinical samples (26) and the prevalence of gene detection varies according to the isolation sources. In Korea, a previous study reported that the *bla*_{OXA} and *bla*_{TEM} genes were detected in 3% and 100% of *Aeromonas* isolates from aquaculture fish [14]. However, a different trend was observed in this study which the *bla*_{OXA} and *bla*_{TEM} genes were detected in *Aeromonas*

isolates from pet turtles that suggest a wide distribution of β -lactamase genes in *Aeromonas* isolates from various sources.

A much higher level of tetracycline resistance was observed amongst aeromonads in our previous study (7) and 78 of tetracycline-resistant isolates were selected to detect their tetracycline resistance determinants (*tetA*, *tetB* and *tetE*). *A. enteropelogenes* and *A. hydrophila* harbored *tetA* and *tetE* genes while other *Aeromonas* species harbored *tetB* and *tetE* genes. Previous reports indicate that the *tetA* and *tetE* determinants are the predominant tetracycline resistance genes in the aquatic environment (16, 27) and both genes code for an efflux pump that eliminates the drug from the cell²⁸. The *tetA*, *tetB* and *tetE* genes are located on the plasmid as well as *tetA* in the transposon (Tn1721) and *tetE* is adjacent to the integrons (15). Han et al. (27) has reported that *tetE* gene was the predominant tetracycline determinant in *Aeromonas* spp. isolated from Korean fish farms and aquariums. However, Kim et al. (29) reported that *tetA* was the most frequent gene in *A. salmonicida* strains isolated from salmonid farms and private aquariums in Korea. The *tetB* gene was detected at a low frequency, while Jacobs and Chenia. (12) reported a lower prevalence of *tetB* genes among *Aeromonas* spp. isolated from the South African aquaculture system.

Detection of virulence encoding genes of *Aeromonas* spp. have been widely applied for evaluating their potential pathogenicity (30, 31). However, the prevalence of virulence-associated genes has rarely been reported in *Aeromonas* strains from pet turtles (7). In the current study, *Aeromonas* isolates were found to possess genes *aerA*, *lip*, *ahyB*, *ser*, *fla*, *gcat* and *ascV*, while genes for DNase (*exu*) was not identified. Especially, none of *A. enteropelogenes* isolates harbored *lip*, *ser*, *exu* and *ascV*

genes. Previous studies have revealed that multiple virulence-associated genes are present in *Aeromonas* isolates and having high heterogeneity in the distribution of virulence-associated genes (10, 30, 31). The pore-forming aerolysin/hemolysin encoded *aer* gene was the most prevalent in this study which was detected in 92% of the total isolates representing all species of the genus. Several studies have reported the high prevalence of the *aer* gene in clinical and environmental *Aeromonas* isolates (30, 32).

The three enterotoxins *act*, *alt*, and *ast* have been implicated as major virulence factors in diarrhoeal disease which had been investigated in our previous study (7). However, the presence of these toxins might not be enough for virulence (31). The temperature-stable metalloprotease with elastolytic activity (*ahyB*) and serine protease (*ser*) play an important role in the invasiveness and establishment of infection (1). In the current study, the *ahyB* and *ser* genes were detected in 59% and 39% of isolates, respectively. None of the *A. enteropelogenes* isolates harbored *ser* gene. The flagella are important appendages for the initial attachment of bacteria to the gastrointestinal epithelium and involve in the subsequent adherence process and biofilm formation (33, 34). The *fla* gene-encoded polar flagella were common among the *Aeromonas* isolates from the aquatic environment. The *fla* gene was detected in 99% of *Aeromonas* isolates from diseased eel in Korea (10). The *gcaT* gene plays a coherent, integrated role in the establishment of pathogenicity of *Aeromonas* spp. by involving in the regulation and secretion of extracellular glycerophospholipid-cholesterol acyltransferase (13). The *gcaT* gene was detected in 68% of *Aeromonas* isolates.

Lipases play a role as virulence factors by interacting with leukocytes or by disturbing several immune system functions through free fatty acids produced by the lipolytic activity. Extracellular lipases secreted by *Aeromonas* spp. actively involve in the alteration of the host plasma membrane and thus increase the severity of infection (35). Among *Aeromonas* strains isolated in the present study, 91% of *A. hydrophila*, 60% of *A. veronii*, 50% of *A. dhakensis* and 50% of *A. caviae* isolates were found to have *lip* gene. Several previous studies reported a high prevalence of *lip* gene among the *Aeromonas* isolates from the aquatic environment (10, 36). Type III secretion system (T3SS) plays a crucial role in host-pathogen interactions by injecting effector toxins directly into the cytosol of host cells (37). The *acsV* gene encodes the T3SS and which was detected in

59% of *Aeromonas* spp. except for *A. enteropelogenes* isolates. The presence of *ascV* gene was previously detected in 68% of *Aeromonas* spp. isolated from diseased farmed fish and farm environment (38). Besides, the high frequency of *ascV* gene was reported in human clinical isolates (37).

The *exu* gene is responsible for DNA hydrolysis which was not detected in this study. The absence of *exu* gene was also reported by Nawaz et al. (13) in *A. veronii* isolated from catfish in the USA. In contrast, the high prevalence of *exu* gene was observed in *Aeromonas* spp. isolated from freshwater lakes in Malaysia (39) and diseased eel in South Korea (10). The specificity of the host or environmental source could be the possible reasons for the absence of *exu* gene in this study.

According to the available literature, this is the first description of these virulence-associated genes in *Aeromonas* of pet turtle origin. Most of *Aeromonas* strains isolated from pet turtles and their environment harboring multiple virulence-associated genes have the potential to be pathogenic. Turtle born aeromonads carrying tetracycline and β -lactams resistance determinants can disseminate through the environment. Collectively, which may pose a public health risk to pet turtle owners, particularly to immunocompromised individuals.

Acknowledgment

This study was supported by the Basic Science Research Program through the National Research Foundation of (KNRF) funded by the Ministry of Education (NRF-2015R1D1A1A01060638) in the Republic of Korea.

Authors declare that no any conflict of interest exists.

References

1. Janda JM, Abbott SL. The genus *Aeromonas*: taxonomy, pathogenicity, and infection. Clin Microbiol Rev 2010; 23: 35–73. <https://doi.org/10.1128/CMR.00039-09>.
2. Kim KT, Kwak D. A case of *Aeromonas hydrophila* infection due to captivity-induced stress in a spectacled caiman (*Caiman crocodilus*). J Anim Plant Sci 2013; 23 (6): 1761–3.
3. Chen J, Zhu N, Kong L, Bei Y, Zheng T, Ding X, He Z. First case of soft shell disease in chinese soft-shelled turtle (*Trionyx sinensis*) associated with *Aeromonas sobria*–*A. veronii* complex. Aquaculture

- 2013; 406–407, 62–7. <https://doi.org/10.1016/j.aquaculture.2013.05.006>.
4. Evangelista-Barreto NS, Vieira RH, Carvalho FCT, et al. *Aeromonas* spp. isolated from oysters (*Crassostrea rhizophorea*) from a natural oyster bed, Ceará, Brazil. *Rev Inst Med Trop São Paulo* 2006; 48: 129–33.
5. Shakir Z, Khan S, Sung K, et al. Molecular characterization of fluoroquinolone-resistant *Aeromonas* spp. isolated from imported shrimp. *Appl Environ Microbiol* 2012; 78: 8137–41. <https://doi.org/10.1128/AEM.02081-12>.
6. Lupescu I, Baraitareanu S. Emerging diseases associated with "New companion animals": review in zoonoses transmitted by reptiles. *Sci Works Ser C Vet Med* 2015; 61: 135–8.
7. Wimalasena SHMP, Shin GW, Hossain S, Heo GJ. Potential enterotoxicity and antimicrobial resistance pattern of *Aeromonas* species isolated from pet turtles and their environment. *J Vet Med Sci* 2017; 79: 921–6. <https://doi.org/10.1292/jvms.16-0493>.
8. Khajanchi BK, Fadl AA, Borchardt MA, et al. Distribution of virulence factors and molecular fingerprinting of *Aeromonas* species isolates from water and clinical samples: suggestive evidence of water-to-human transmission. *Appl Environ Microbiol* 2010; 76: 2313–25. <https://doi.org/10.1128/AEM.02535-09>.
9. Ottaviani D, Parlani C, Citterio B, et al. Putative virulence properties of *Aeromonas* strains isolated from food, environmental and clinical sources in Italy: a comparative study. *Int J Food Microbiol* 2011; 144: 538–45. <https://doi.org/10.1016/j.ijfoodmicro.2010.11.020>.
10. Yi SW, You MJ, Cho HS, Lee CS, Kwon JK, Shin GW. Molecular characterization of *Aeromonas* species isolated from farmed eels (*Anguilla japonica*). *Vet Microbiol* 2013; 164: 195–200. <https://doi.org/10.1016/j.vetmic.2013.02.006>.
11. Deng YT, Wu Y, Tan AP, et al. Analysis of antimicrobial resistance genes in *Aeromonas* spp. isolated from cultured freshwater animals in China. *Microb Drug Resist* 2014; 20: 350–6. <https://doi.org/10.1089/mdr.2013.0068>.
12. Jacobs L, Chenia HY. Characterization of integrons and tetracycline resistance determinants in *Aeromonas* spp. isolated from South African aquaculture systems. *Int J Food Microbiol* 2007; 114: 295–306. <https://doi.org/10.1016/j.ijfoodmicro.2006.09.030>.
13. Nawaz M, Khan SA, Khan AA, et al. Detection and characterization of virulence genes and integrons in *Aeromonas veronii* isolated from catfish. *Food Microbiol* 2010; 27: 327–31. <https://doi.org/10.1016/j.fm.2009.11.007>.
14. Yi SW, Chung TH, Joh SJ, Park C, Park BY, Shin GW. High prevalence of bla_{CTX-M} group genes in *Aeromonas dhakensis* isolated from aquaculture fish species in South Korea. *J Vet Med Sci* 2014; 76: 1589–93. <https://doi.org/10.1292/jvms.14-0274>.
15. Akinbowale Ol, Peng H, Barton MD. Diversity of tetracycline resistance genes in bacteria from aquaculture sources in Australia. *J Appl Microbiol* 2007; 103: 2016–25. <https://doi.org/10.1111/j.1365-2672.2007.03445.x>.
16. Nawaz M, Sung K, Khan SA, Khan AA, Steele R. Biochemical and molecular characterization of tetracycline-resistant *Aeromonas veronii* isolates from catfish. *Appl Environ Microbiol* 2006; 72: 6461–6. <https://doi.org/10.1128/AEM.00271-06>.
17. Henriques IS, Fonseca F, Alves A, Saavedra MJ, Correia A. Occurrence and diversity of integrons and β -Lactamase genes among ampicillin-resistant isolates from estuarine waters. *Res Microbiol* 2006; 157: 938–47. <https://doi.org/10.1016/j.resmic.2006.09.003>.
18. Maravić A, Skočibušić M, Šamanić I, et al. *Aeromonas* spp. simultaneously harboring bla(CTX-M-15), bla(SHV-12), bla(PER-1) and bla(FOX-2), in wild-growing mediterranean mussel (*Mytilus galloprovincialis*) from Adriatic sea, Croatia. *Int J Food Microbiol* 2013; 166: 301–8. <https://doi.org/10.1016/j.ijfoodmicro.2013.07.010>.
19. Tacão M, Correia A, Henriques I. Resistance to broad-spectrum antibiotics in aquatic systems: anthropogenic activities modulate the dissemination of blaCTX-M-like genes. *Appl Environ Microbiol* 2012; 78: 4134–40. <https://doi.org/10.1128/AEM.00359-12>.
20. Piotrowska M, Popowska M. The Prevalence of antibiotic resistance genes among *Aeromonas* species in aquatic environments. *Ann Microbiol* 2014; 64: 921–34. <https://doi.org/10.1007/s13213-014-0911-2>.
21. Wimalasena SHMP, De Silva BCJ, Hossain S, Pathirana HNKS, Heo GJ. Prevalence and characterization of quinolone resistance genes in *Aeromonas* species isolated from pet turtle in Korea. *J Glob Antimicrob Resist* 2017; 11: 34–8 <https://doi.org/10.1016/j.jgar.2017.06.001>.
22. Chung T, Yi S, Kim B, Kim W, Shin GW. Identification and antibiotic resistance profiling of bacterial isolates from septicemic soft-shelled turtles

(*Pelodiscus Sinensis*). Vet Med 2017; 62: 169–77. <https://doi.org/10.17221/65/2016-VETMED>.

23. Bhaskar M, Dinoop KP, Mandal J. Characterization of ceftriaxone-resistant *Aeromonas* spp. isolates from stool samples of both children and adults in southern India. J Health Popul Nutr 2015; 33: e26. <https://doi.org/10.1186/s41043-015-0036-7>.

24. Figueira V, Vaz-Moreira I, Silva M, Manaia CM. Diversity and antibiotic resistance of *Aeromonas* spp. in drinking and waste water treatment plants. Water Res 2011; 45: 5599–611. <https://doi.org/10.1016/j.watres.2011.08.021>.

25. Piotrowska M, Przygodzińska D, Matyjewicz K, Popowska M. Occurrence and variety of β -lactamase genes among *Aeromonas* spp. isolated from urban wastewater treatment plant. Front Microbiol 2017; 8: e863 <https://doi.org/10.3389/fmicb.2017.00863>.

26. Puah SM, Puthucheary SD, Liew FY, Chua KH. *Aeromonas* aquariorum clinical isolates: antimicrobial profiles, plasmids and genetic determinants. Int J Antimicrob Agents 2013; 41: 281–4. <https://doi.org/10.1016/j.ijantimicag.2012.11.012>.

27. Han JE, Kim JH, Choresca CH, et al. Prevalence of *tet* gene and complete genome sequencing of *tet* gene-encoded plasmid (PAHH01) isolated from *Aeromonas* species in South Korea. J Appl Microbiol 2012; 112: 631–8. <https://doi.org/10.1111/j.1365-2672.2012.05237.x>.

28. Chopra I, Roberts M. Tetracycline antibiotics: mode of action, applications, molecular biology, and epidemiology of bacterial resistance. Microbiol Mol Biol Rev 2001; 65: 232–60. <https://doi.org/10.1128/MMBR.65.2.232-260.2001>.

29. Kim JH, Hwang SY, Son JS, et al. Molecular characterization of tetracycline- and quinolone-resistant *Aeromonas salmonicida* isolated in Korea. J Vet Sci 2011; 12: 41–8. <https://doi.org/10.4142/jvs.2011.12.1.41>.

30. Chacón MR, Figueras MJ, Castro-Escarpulli G, Soler L, Guarro J. Distribution of virulence genes in clinical and environmental isolates of *Aeromonas* spp. Antonie Van Leeuwenhoek 2003; 84: 269–78.

31. Sha J. Role of various enterotoxins in *Aeromonas hydrophila*-induced gastroenteritis: Generation of enterotoxin gene-deficient mutants and evaluation of their enterotoxic activity. Infect Immun 2002; 70: 1924–35. <https://doi.org/10.1128/IAI.70.4.1924-1935.2002>.

32. Ghenghesh KS, Ahmed SF, Cappuccinelli P, Klena JD. Genospecies and virulence factors of *Aeromonas* species in different sources in a north African country. Libyan J Med 2014; 9 (1): e25497. <https://doi.org/10.3402/ljm.v9.25497>.

33. Kirov SM, Castrisios M, Shaw JG. *Aeromonas* flagella (polar and lateral) are enterocyte adhesins that contribute to biofilm formation on surfaces. Infect Immun 2004; 72: 1939–45. <https://doi.org/10.1128/IAI.72.4.1939-1945.2004>.

34. Umelo E. Identification and molecular characterization of two tandemly located flagellin genes from *Aeromonas salmonicida* A449. J Bacteriol 1997; 179: 5292–9.

35. Pemberton JM, Kidd SP, Schmidt R. Secreted enzymes of *Aeromonas*. FEMS Microbiol Lett 1997; 152: 1–10. <https://doi.org/10.1111/j.1574-6968.1997.tb10401.x>.

36. Sen K, Rodgers M. Distribution of six virulence factors in *Aeromonas* species isolated from US drinking water utilities: A PCR identification. J Appl Microbiol 2004; 97: 1077–86. <https://doi.org/10.1111/j.1365-2672.2004.02398.x>.

37. Vilches S, Urgell C, Merino S, et al. Complete type III secretion system of a mesophilic *Aeromonas hydrophila* strain. Appl Environ Microbiol 2004; 70: 6914–9. <https://doi.org/10.1128/AEM.70.11.6914-6919.2004>.

38. Carvalho-Castro GA, Lopes CO, Leal CAG, Cardoso PG, Leite RC, Figueiredo HCP. Detection of type III secretion system genes in *Aeromonas hydrophila* and their relationship with virulence in Nile tilapia. Vet Microbiol 2010; 144: 371–6. <https://doi.org/10.1016/j.vetmic.2010.01.021>.

39. Khor WC, Puah SM, Tan JAMA, Puthucheary S, Chua KH. Phenotypic and genetic diversity of *Aeromonas* species isolated from fresh water lakes in Malaysia. PLOS ONE 2015; 10 (12): e0145933. <https://doi.org/10.1371/journal.pone.0145933>.

PRISOTNOST DETERMINANT ZA DOLOČITEV DOMNEVNE VIRULENCE TER GENOV ZA ODPORNOST NA TETRACIKLIN IN β -LAKTAM VRST *Aeromonas* IZOLIRANIH IZ LJUBITELJSKIH VRST ŽELV IN IZ NJIHOVEGA OKOLJA

S.H.M.P.Wimalasena, G-J. Heo

Izvleček: Namen študije je bil določiti bakterije *Aeromonas* spp., izolirane iz desetih priljubljenih vrst hišnih želv in njihovega okolja, z namenom ocenjevanja potencialnega tveganja hišnih želv kot vira genov, povezanih z virulenco, ter determinante odpornosti proti tetraciklinom in β -laktamom. Prisotnost osmih virulentnih genov (*ser*, *aer*, *exu*, *lip*, *fla*, *ascV*, *ahyB* in *gcat*) ter genov za odpornost na tetracikline (*tetA*, *tetB* in *tetE*) in β -laktame (*bla*_{TEM}, *bla*_{SHV}, *bla*_{OXA} in *bla*_{CTX-M}) je bila ocenjena s konvencionalnimi testi PCR. Najbolj pogost je bil Gen *aerA* (92 %), sledili so geni *fla* (75 %), *gcat* (68 %), *ahyB* (59 %), *ser* (39 %), *lip* (37 %) in *ascV* (25 %). Nobeden od izolatov ni imel pomnoženega gena *exu*, povezanega z DNAzo. *A. hydrophila*, *A. dharkensis*, *A. veronii* in *A. caviae* so vsebovali sedem testiranih genov virulence, razen *exu*, medtem ko so bili v *A. enteropelogenih* odkriti le štiri virulentni geni. Med 75 izolati, odpornimi na tetracikline, so bili geni *tetA*, *tetE* in *tetB* odkriti v 38, 26 oziroma 6 izolatih. Med preizkušeni genoma za odpornost proti β -laktamu so bili geni *bla*_{OXA} in *bla*_{TEM} odkriti pri 54 % oziroma 36 % izolatov, odpornih proti β -laktamu.

V nobenem vzorcu nista bila zaznana gena *bla*_{CTX-M} in *bla*_{SHV}. Rezultati študije kažejo, da bakterije *Aeromonas* spp. iz hišnih želv lahko imajo potencialne virulentne gene in gene za odpornost proti tetraciklinu in β -laktamom, in lahko potencialno ogrožajo zdravje lastnikov hišnih želv, zlasti imunsko oslabljenih posameznikov.

Ključne besede: *Aeromonas* spp.; geni povezani z virulenco; odpornost na tetracikline; rezistenca na β -laktami; ljubiteljske vrste želv